

# Impact of Inhibition of the TCA cycle on ATP and NADH levels in *Escherichia coli* during Bacterial Replication

Raina De La Cruz, Briana Mercado, Aiswarya Suriaprakash, and Jessica O'Hara

## EXPANSION OF RESEARCH


- Quantify ATP, NADH, and glucose levels before and after cell lysis to further understand the impacts of each knockout on cellular metabolism and the resulting impact on T4 replication.
- Expand range of genes in the TCA cycle to manipulate (*gltA*, *maeA*, *acnB*, *sucC*)
  - This will be quantified by utilizing a plate reader (@600nm) in both growth and lysis curves over eight hours.
  - The growth curve will measure the growth of the new genes compared to wildtype (WT) *E. coli*
  - The double agar overlay method will be utilized to grow and analyze T4 and T4r bacteriophage growth with the wild type and gene knockout of choice.

Gene	TCA Cycle Enzyme
<i>gltA</i>	Citrate Synthase
<i>maeA</i>	Pyruvate Dehydrogenase
<i>acnB</i>	Aconitase
<i>sucC</i>	Succinyl-CoA Synthetase


**Objective:** Understand ways to treat bacterial diseases in the face of antibiotic resistance, specifically phage therapy.

## INTRODUCTION TO *FUM* AND *ICD* GENES

The tricarboxylic acid (TCA) is the metabolic pathway that both genes of interest are part of. The TCA cycle consists of a series of reactions that occur within aerobic respiration. This is not the most ATP generative part of aerobic respiration; rather, it creates high energy carriers, NADH and FADH<sub>2</sub>. The *fumA* gene is one of the three fumarase genes that creates the enzyme need for the conversion of fumarate to S-malate, in the last few processes of the TCA cycle.




**Figure 2.** *Δicd* Reaction in Krebs Cycle. Diagram of the TCA cycle and gene knockout implications.



**Figure 1.** *ΔfumA* in the TCA cycle Illustration of the TCA cycle and gene knockout implications.


The *icd* gene connects the TCA cycle to the glyoxylate bypass pathway. It is the first enzyme in a bacterial pathway to be regulated by phosphorylation.

## PLAQUE ASSAYS SHOW BACTERIAL LYSIS



**Figure 3.** *ΔfumA* plaque assays

A double agar overlay method was used with T4 bacteriophage in the wild type (A and C) and *ΔfumA E. coli* strains (B and D) at dilutions of 10<sup>-6</sup> (A and B) and 10<sup>-7</sup> (C and D). There are less PFU in the *ΔfumA* strain than the wild type.



**Figure 4.** *Δicd* plaque assays

The plaque assays feature plaques of similar size and shape, indicating similar lysis of bacterial cells by phage.

## ΔFUMA AND ΔICD LED TO DELAYED BACTERIOPHAGE REPLICATION


**Figure 5. Growth Curves**

A) Growth of *Δicd* and *ΔfumA* in LB Media

Growth of both knockout strains were not impacted in LB media.

B) Growth of *Δicd* and *ΔfumA* in M9 Media

Growth of the *Δicd* strain was significantly less in M9 media.




**Figure 6. Lysis Curves**

A) *ΔfumA*

Lysis of the parent and *ΔfumA* strain with T4 phage was observed over time by measuring absorbance at 600nm. The *ΔfumA* strain experienced slower lysis by phage than the parent strain.

B) *Δicd*

Lysis of the parent and *Δicd* strain with T4r phage was observed over time by measuring absorbance at 600nm. The *Δicd* strain experienced slower lysis by phage than the parent strain



## REFERENCES & ACKNOWLEDGEMENTS

- ATP assay Kit (Colorimetric/Fluorometric) (ab83355). (2021, April 12). Retrieved April 16, 2021, from <https://www.abcam.com/atp-assay-kit-colorimetricfluorometric-ab83355.html>
- "Escherichia Coli K-12 Substr. MG1655 Icd." <https://biocyc.org/gene?orgid=ECOLI&id=EG10489> (April 14, 2021).
- Gerdes SY, et. al. Experimental determination and system level analysis of essential genes in Escherichia coli MG1655. *J Bacteriol.* 2003 Oct;185(19):5673-84.
- Glucose (HK) Assay Kit GAHK20*. Sigma-Aldrich. Retrieved April 16, 2021, from <https://www.sigmaaldrich.com/catalog/product/sigma/gahk20>
- Kabir, M.M., and Shimizu, K. (2004). Metabolic regulation analysis of icd-gene knockout Escherichia coli based on 2D electrophoresis with MALDI-TOF mass spectrometry and enzyme activity measurements. *Appl Microbiol Biotechnol* 65, 84–96.
- Leonardo, M R, Y Dailly, and D P Clark. 1996. "Role of NAD in Regulating the AdhE Gene of Escherichia Coli." *Journal of bacteriology* 178(20): 6013–18.
- Li, J., Li, Y., Cui, Z. *et al.* Enhancement of succinate yield by manipulating NADH/NAD<sup>+</sup> ratio and ATP generation. *Appl Microbiol Biotechnol* 101, 3153–3161 (2017). <https://doi.org/10.1007/s00253-017-8127-6>
- M R Leonardo, Y Dailly, D P Clark *Journal of Bacteriology* Oct 1996, 178 (20) 6013-6018; DOI: 10.1128/jb.178.20.6013-6018.1996

Financial support for this project was provided by the First-Year Innovation and Research Experience at the University of Maryland, College Park.

## ΔICD FOUND TO DECREASE ATP LEVELS IN *E. COLI*

- Deletion of the *icd* gene led to reduced specific growth rate and reduced specific glucose consumption rate because of the lower intracellular ATP/ADP and NADPH/NAD<sup>+</sup> ratios compared to the parent strain<sup>5</sup>.
- Since *ΔfumA* followed the same trends in growth and lysis curves, it is assumed that the quantification of ATP/ADP and NADPH/NAD<sup>+</sup> ratios in this strain and other TCA cycle enzyme knockout strains will provide similar results.

## QUANTIFICATION METHODS

These assays will be performed to quantify NAD-NADH ratios, glucose consumption rate, and ATP levels during growth and viral replication.

NAD-NADH Pool Isolation and NAD Cycling Assay <sup>6</sup>	Sigma-Aldrich Glucose (HK) Assay Kit <sup>4</sup>	ATP Assay Kit (Colorimetric) <sup>1</sup>
<ol style="list-style-type: none"><li>Samples will be centrifuged and the subsequent supernatant will be removed and will immediately be frozen in a dry ice-ethanol bath.</li><li>HCl (for NAD extraction) or NaOH (for NADH extraction) is added to the pellets.</li><li>Cycling assays will be performed for both, the reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-di phenyl tetrazolium bromide (MTT) is directly proportional to levels of NAD and NADH @ 570 nm.</li></ol>	<ol style="list-style-type: none"><li>Glucose is phosphorylated by ATP with the help of hexokinase → Glucose-6-Phosphate (G6P) and ADP are products.</li><li>G6P is oxidized to 6-phosphogluconate in the presence of NAD in a reaction catalyzed by glucose-6-phosphate dehydrogenase (G6PDH)</li><li>During the oxidation, an equimolar amount of NAD is reduced to NADH.</li><li>The resulting increase in absorbance @ 340 nm is directly proportional to the concentration of glucose.</li><li>Glucose consumption rate is measured by measuring glucose concentration (intra or extracellular) over time.</li></ol>	<ol style="list-style-type: none"><li>ATP is made through metabolic reactions in the cells → ATP is then used to phosphorylate glycerol.</li><li>The phosphorylation of glycerol reaction produces byproduct (Resorufin) → byproduct is then stained red via assay and able to be quantified by optical density (OD) readings @ 570 nm.</li></ol>